

Cystatin C (E.coli) Human, Rabbit Polyclonal Antibody

Product Data Sheet

Source of Antigen: *E. coli*

Host: Rabbit

Cat. No.:

RD181009100

(0.1 mg)

Other names: Post G-globulin, Cystatin-3, Neuroendocrine basic polypeptide, Gamma-trace, Post-gamma-globulin, CST3

Research topic

Neural tissue markers, Renal disease

Preparation

The antibody was raised in rabbits by immunization with the recombinant Human Cystatin C.

Amino Acid Sequence

The immunization antigen (14.5 kDa) is a protein containing 129 AA of recombinant Human Cystatin C. N-terminal His-tag, 9 extra AA (highlighted).

MKHHHHHHAS SP-GKPPRLVG GPMDASVEEE GRRALDFAV GEYNKASNDM YHSRALQVVR ARKQIVAGVN YFLDVELGRT
TCTKTQPNLD NCPFHDQPHL KRKAFCSFQI YAVPWQGTMT LSKSTCQDA

Species Reactivity

Human

Not yet tested in other species.

Purification Method

Immunoaffinity chromatography on a column with immobilized recombinant Human Cystatin C.

Antibody Content

0.1 mg (determined by BCA method, BSA was used as a standard)

Formulation

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. **AZIDE FREE.**

Reconstitution

Add 0.1 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

Shipping

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage/Stability

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80°C. Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C.

Expiration

See vial label.

Lot Number

See vial label.

Quality Control Test


Indirect ELISA - to determine titer of the antibody

SDS PAGE - to determine purity of the antibody

Applications

ELISA, Immunohistochemistry, Western blotting

Antibodies application



Introduction to the Molecule

Cystatin C is a non-glycosylated basic single-chain protein consisting of 120 amino acids with a molecular weight of 13.36 kDa and is characterized by two disulfide bonds in the carboxy-terminal region. It belongs to the cystatins superfamily which inactivates lysosomal cysteine proteinases, e.g. cathepsin B, H, K, L and S. Imbalance between Cystatin C and cysteine proteinases is associated with inflammation, renal failure, cancer, Alzheimer's disease, multiple sclerosis and hereditary Cystatin C amyloid angiopathy. Its increased level has been found in patients with autoimmune diseases, with colorectal tumors and in patients on dialysis. Serum Cystatin C seems to be better marker of glomerular filtration rate than creatinine. On the other hand, low concentration of Cystatin C presents a risk factor for secondary cardiovascular events.

HEADQUARTERS: BioVendor Laboratorní medicína, a.s.	Karasek 1767/1	621 00 Brno CZECH REPUBLIC	Phone: +420-549-124-185 Fax: +420-549-211-460	E-mail: info@biovendor.com sales@biovendor.com Web: www.biovendor.com
AUSTRIA: BioVendor GesmbH	Gaudenzdorfer Gürtel 43-45	1120 Vienna AUSTRIA	Phone: +43-1-89090-25 Fax: +43-1-89051-63	E-mail: infoAustria@biovendor.com
GERMANY, SWITZERLAND: BioVendor GmbH	Otto-Hahn-Straße 16	34123 Kassel GERMANY	Phone: +49-6221-433-9100 Fax: +49-6221-433-9111	E-mail: infoEU@biovendor.com
USA, CANADA AND MEXICO: BioVendor LLC	128 Bingham Rd. Suite 1300	Asheville, NC 28806 USA	Phone: +1-828-575-9250 +1-800-404-7807 Fax: +1-828-575-9251	E-mail: infoUSA@biovendor.com